Biomedical Optics EXPRESS

Moxifloxacin based fluorescence imaging of intestinal goblet cells: supplement

SEUNGHUN LEE,^{1,6} SEONGHAN KIM,^{1,6} KWANGWOO NAM,² SUN YOUNG KIM,³ SEUNGRAG LEE,⁴ SEUNG-JAE MYUNG,^{3,7} AND KI HEAN KIM^{1,5,8}

This supplement published with The Optical Society on 23 Sepember 2020 by The Authors under the terms of the Creative Commons Attribution 4.0 License in the format provided by the authors and unedited. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Supplement DOI: https://doi.org/10.6084/m9.figshare.12971684

Parent Article DOI: https://doi.org/10.1364/BOE.402350

¹Department of Mechanical Engineering, Pohang University of Science and Technology, 77 Cheongam-ro, Nam-gu, Pohang, Gyeongbuk 37673, South Korea

²Department of Internal Medicine, Dankook University College of Medicine, 201 Manghyang-ro, Dongnam-gu, Cheonan, Chungnam 31116, South Korea

³Department of Gastroenterology, Digestive Diseases Research Center, Asan Institute for Life Sciences, Asan Medical Center, University of Ulsan College of Medicine, 88 Olympic-ro, 43-gil, Songpa-gu, Seoul 05505, South Korea

⁴Medical Device Development Center, Osong Medical Innovation Foundation, 123 Osongsaengmyeong-ro, Heungdeok-gu, Cheongju, Chungbuk 28160, South Korea

⁵Division of Integrative Biosciences and Biotechnology, Pohang University of Science and Technology, 77 Cheongam-ro, Nam-gu, Pohang, Gyeongbuk 37673, South Korea

⁶These authors contributed equally to this work

⁷sjmyung@amc.seoul.kr

⁸kiheankim@postech.ac.kr

Moxifloxacin based fluorescence imaging of intestinal goblet cells




Fig. S1. Autofluorescence emission spectrum of the confocal miniprobe (Gastroflex UHD, Mauna Kea Technology) at the excitation wavelength of 405 nm.




Fig. S2. Signal to background ratio analysis of moxifloxacin based intestinal goblet cell imaging in the normal mouse colon.